



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A SINGULAR VALUE OF π .

BY PROF. J. W. NICHOLSON, LOUISIANA STATE UNIV., BATON ROUGE, LA.

ON page 291 of Ray's Calculus may be seen a demonstration of the following well known theorem of Wallis :

$$\frac{\pi}{2} = \frac{2 \cdot 2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \cdot 8 \cdot 8 \dots}{1 \cdot 3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 9 \dots} \quad (1)$$

By the binomial formula

$$(1-1)^n = 1 - n + \frac{n(n-1)}{2} - \frac{n(n-1)(n-2)}{2 \cdot 3} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 3 \cdot 4} \dots \quad (2)$$

Factoring,

$$(1-1)^n = \frac{(1-n)(2-n)(3-n)(4-n) \dots}{1 \cdot 2 \cdot 3 \cdot 4 \dots} \quad (3)$$

Substituting $-n$ for n ,

$$(1-1)^{-n} = \frac{(1+n)(2+n)(3+n)(4+n) \dots}{1 \cdot 2 \cdot 3 \cdot 4 \dots} \quad (4)$$

Multiplying (3) by (4),

$$(1+1)^n(1-1)^{-n} = \frac{(1-n^2)(4-n^2)(9-n^2)(16-n^2) \dots}{1 \cdot 2 \cdot 2 \cdot 3 \cdot 3 \cdot 4 \cdot 4 \dots} \quad (5)$$

Substituting $\frac{1}{2}$ for n , and reducing,

$$(1-1)^{\frac{1}{2}}(1-1)^{-\frac{1}{2}} = \frac{1 \cdot 3 \cdot 3 \cdot 5 \cdot 5 \cdot 7 \cdot 7 \cdot 9 \dots}{2 \cdot 2 \cdot 4 \cdot 4 \cdot 6 \cdot 6 \cdot 8 \cdot 8 \dots} \quad (6)$$

Combining (1) and (6),

$$\pi = \frac{2}{(1-1)^{\frac{1}{2}}(1-1)^{-\frac{1}{2}}}.$$

ANSWER TO PROF. SCHEFFER'S QUER (P. 31, VOL. VIII.)

BY C. B. SEYMOUR, ATTORNEY AT LAW, LOUISVILLE, KY.

Query.—"If of any curve we find the evolute, and of the latter the evolute, and so on ad infin., the ultimate evolute is a cycloid. How is this proved?"

Answer.—The proposition stated is not correct.

Let s_0 be the length of the given curve, measured from the origin to any point (the origin being a point on the curve). Let β_0 be the inclination of the tangent at that point to the axis of abscissas, and let R_0 be the radius of curvature at that point. Let s_n, β_n, R_n be like quantities for the corres-